

1 **Modeling fuel-, vehicle type-, and age-specific CO₂** 2 **emissions from global on-road vehicles, 1970-2020**

3 Liu Yan¹, Qiang Zhang², Bo Zheng³, Kebin He¹

4 ¹ State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment,
5 Tsinghua University, Beijing, People's Republic of China

6 ² Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System
7 Science, Tsinghua University, Beijing, People's Republic of China

8 ³ Institute of Environment and Ecology, Tsinghua Shenzhen International Graduate School, Tsinghua
9 University, Shenzhen, China

10 *Correspondence to:* Qiang Zhang (qiangzhang@tsinghua.edu.cn)

11 **Abstract.** Vehicles are among the most important contributors to global anthropogenic CO₂ emissions.
12 However, the lack of fuel-, vehicle type-, and age-specific information about global on-road CO₂
13 emissions in existing datasets, which are available only at the sector level, makes these datasets
14 insufficient to support the establishment of emission mitigation strategies. Thus, a fleet turnover model
15 is developed in this study, and CO₂ emissions from global on-road vehicles from 1970 to 2020 are
16 estimated for each country. Here, we analyze the evolution of the global vehicle stock over 50 years,
17 identify the dominant emission contributors by vehicle and fuel type, and further characterize the age
18 distribution of on-road CO₂ emissions. We find that trucks accounted for less than 5% of global vehicle
19 ownership but represented more than 20% of on-road CO₂ emissions in 2020. The contribution of diesel
20 vehicles to global on-road CO₂ emissions doubled during the 1970-2020 period, driven by the shift in
21 the fuel-type distribution of vehicle ownership. The proportion of CO₂ emissions from vehicles in
22 developing countries such as China and India in terms of global emissions from newly registered vehicles
23 significantly increased after 2000, but global CO₂ emissions from vehicles that survived more than 15
24 years in 2020 still originated mainly from developed countries such as the United States and countries in
25 the European Union.

1 Introduction

To meet the Paris Agreement's 1.5°C long-term temperature goal, many efforts have been made to determine pathways for reducing the emissions of greenhouse gases such as CO₂ (Matthews & Caldeira, 2008; Meinshausen et al., 2009; Rogelj et al., 2018; Davis et al., 2018). Historical emission data and consistent emission series of on-road vehicles, which are key sources of CO₂ emissions, are important inputs for Earth system models, atmospheric chemistry and transport models, and integrated assessment models to support studies on both climate change and global climate governance (Bhalla et al., 2014; Janssens-Maenhout et al., 2019; Lelieveld et al., 2015; Niklas et al., 2020; Shindell et al., 2011; Silva et al., 2016; Unger et al., 2010). Thus, estimating long-term CO₂ emissions from global on-road vehicles with detailed source information is necessary as deep greenhouse gas emission reductions are pursued.

Several global emission inventories that cover emissions from on-road vehicles have been developed and are widely used in global research and modeling. CO₂ emissions from on-road vehicles can be derived from global anthropogenic emission inventories, including the Emissions Database for Global Atmospheric Research (EDGAR), the Open-source Data Inventory for Atmospheric CO₂ (ODIAC), the Carbon Emission and Accounts Datasets (CEADs), and the Peking University (PKU)-CO₂ inventory. On-road CO₂ emissions are estimated with the total fuel consumption of the road sector at the country level and fleet average emission factors in EDGAR (Amstel et al., 1999; Crippa et al., 2016; Crippa et al., 2018; Janssens-Maenhout et al., 2019). Following the method in EDGAR, local data sources are introduced more often in ODIAC (Boden et al., 2016; Boden et al., 2017; Od et al., 2018), CEDS (Hoesly et al., 2018) and PKU-CO₂ (Wang et al., 2013) when estimating on-road CO₂ emissions. Global CO₂ emissions from on-road vehicles in these widely used emission inventories are estimated as a whole at the sector level in each country using the fuel-based method, and fleet structure information (e.g., fuel-, vehicle type-, and age-specific characteristics) on on-road CO₂ emissions is omitted. Technology-based models such as the Greenhouse Gas and Air Pollution Interactions and Synergies (GAINS) (Klimont et al., 2017) and Speciated Pollutant Emissions Wizard (SPEW)-Trend (Tami et al., 2004 and 2007; Yan et al., 2011 and 2014) models can be used to describe fleet structure information on emissions from global on-road vehicles, but emission inventories built on these models include only emissions of air pollutants.

Here, a new global inventory of fuel-, vehicle type-, and age-specific CO₂ emissions from on-road vehicles for each country from 1970 to 2020 is developed with the global fleet turnover model, in which

six types of fuel, five types of vehicles, and 231 countries are considered. Based on this inventory, we analyze the evolution of the global vehicle stock over 50 years; identify the dominant emission contributors by vehicle and fuel type; and further characterize the age distribution of on-road CO₂ emissions. Compared to the publicly available on-road CO₂ emissions from previous studies, CO₂ emissions in this study have more detailed source categories which are refined into vehicle and fuel type. And with the age distribution simulated by our fleet turnover model, CO₂ emissions offered in this study would better support the policy-making of emission mitigation.

2 Materials and methods

2.1 Methodological framework

For a given country c , the annual CO₂ emissions from on-road vehicles in year y are estimated as follows:

$$Emis_{c,y,v,f} = \sum_{i=0}^{i=T} Stock_{c,y,v} \times X_{c,y,v,i} \times FuelR_{c,y,v,f} \times VKT_{c,y,v,f} \times FE_{c,y,v,f} \times EF_{c,f}, \quad (1)$$

$$Stock_{c,y,v} = V_{c,y,v}^* \times e^{\alpha_{c,v}} e^{\beta_{c,v} E_{c,y}} \times Population_{c,y}, \quad (2)$$

$$Stock_{c,y,v} = \sum_{i=0}^{i=T} Sale_{c,y-i,v} \times Surv_{c,v,i}, \quad (3)$$

$$X_{c,y,v,i} = Sale_{c,y-i,v} \times Surv_{c,v,i} / \sum_{i=0}^{i=T} Sale_{c,y-i,v} \times Surv_{c,v,i}, \quad (4)$$

$$Fuel_{c,y,f} = \sum_v Stock_{c,y,v} \times FuelR_{c,y,v,f} \times VKT_{c,y,v,f} \times FE_{c,y,v,f}, \quad (5)$$

where y is the target year, which ranges from 1970 to 2020; i is the age of the vehicles registered in year $(y - i)$; T is the lifetime of vehicles; v is the vehicle type, which includes two types of light-duty vehicles, namely, passenger cars (PLDVs) and light commercial vehicles (CLDVs), two types of heavy-duty vehicles, namely, buses and trucks, and motorcycles (MCs); and f is the fuel type, which includes gasoline, diesel, natural gas (NG), liquefied petroleum gas (LPG), electricity, and other fuels. As shown in Equation 1, annual CO₂ emissions ($Emis_{c,y,v,f}$) are estimated by the vehicle stock ($Stock_{c,y,v}$), the fleet-average fuel structure ($FuelR_{c,y,v,f}$), the annual average kilometers traveled ($VKT_{c,y,v,f}$), the fleet-average fuel economy ($FE_{c,y,v,f}$), the age distribution of the vehicle stock ($X_{c,y,v,i}$), and the CO₂ emission factor ($EF_{c,f}$). $Stock_{c,y,v}$ can be modeled using the Gompertz function (Equation 2), which is an S-shaped curve determined by two negative parameters (α and β), with the saturated

vehicle stock per 1000 people (V^*), per capita GDP (E), and population ($Population_{c,y}$) as inputs. The age distribution of the vehicle stock ($X_{c,y,v,i}$), which represents the proportion of surviving vehicles registered in year $(y - i)$ in target year y , is modeled on the basis of the dynamic balance function (Equation 3 and 4) using the number of newly registered vehicles ($Sale_{c,y-i,v}$) and survival rates ($Surv_{c,v,i}$). Fuel consumption by vehicle type, which is calculated using $Stock_{c,y,v}$, $X_{c,y,v,i}$, $FuelR_{c,y,v,f}$, $VKT_{c,y,v,f}$, and $FE_{c,y,v,f}$, is constrained by total on-road fuel consumption ($Fuel_{c,y,f}$) at the country level (Equation 5).

In this study, the fleet turnover emission model (Figure 1) is constructed based on equations 1-5. Specifically, we first build an integrated vehicle stock database by combining and harmonizing the available vehicle stock data from a series of global, regional and national statistics and filling data gaps with the modeled stock based on the Gompertz function (Equation 2). Second, the age distribution of the stock is simulated with a combined vehicle sale statistical database and an integrated vehicle stock database using the dynamic balance function (Equation 3 and 4). Third, vehicular fuel consumption is estimated using outputs from the first two steps and other vehicle activity-related data and is constrained by national fuel consumption statistics (Equation 5). Finally, fuel- and vehicle type-specific CO_2 emissions from global on-road vehicles from 1970 to 2020 are modeled on the basis of constrained vehicular fuel consumption and CO_2 emission factors (Equation 1).




Figure 1: Schematic methodology for estimating vehicular CO₂ emissions.

2.2 Modeling the vehicle stock

In the first step, an integrated vehicle stock database from 1970 to 2020 was constructed with both statistical and modeled data. The statistical data used in this study was collected from various available vehicle stock statistics, in which global statistics were used as the default vehicle stock and local statistics were used to supplement and amend the default data. When statistical data was unavailable for a country in a given year, vehicle stock modeled by the Gompertz function was used.

To determine the default vehicle stock database, two widely used vehicle stock statistics from the World Road Statistics (WRS) 2021 Edition (IRF) and the International Organization of Motor Vehicle Manufacturers (OICA) were collected and compared. We found that the trends of vehicle stock in the WRS and OICA data were similar, but the absolute value of the vehicle stock in the OICA data was lower than that in the WRS data, especially for developing countries (Figure S2). Taking India as an example, the vehicle stock in the OICA data was 85% less than that in the WRS data. To further confirm the reliability of these two global databases, local statistics were used for comparison. The WRS data were more similar to the local vehicle statistics than were the OICA data (Figure S2). After comprehensive consideration of spatiotemporal coverage, updating frequency and stability, and data reliability, the WRS data were used as the default for global vehicle statistics, and the OICA data were used if there were no data available from the WRS.

We also collected a series of local statistics as supplements and amendments to the global vehicle statistics, in which 49 developing and developed countries were included (ACEA; CEIC; EC; JAMA; MEIC; MOSPI; NBS; TEDB). By coupling multiple global and local vehicle databases, a combined vehicle statistical database by vehicle category was established in this study. As the division of vehicle types varied among statistics, we established a mapping relationship of vehicle types between this study and other data sources (Table S2).

Given that statistical data of vehicle was unavailable before 2000 for most countries, the Gompertz function, which was often applied to establish the relationship between vehicle ownership and an economic indicator (Dargay and Gately, 1999; Dargay et al., 2007; Huo and Wang, 2012), was subsequently used in this study to model the vehicle stock. In this study, per capita GDP was calculated with national GDP (NBS; UNdata; WB) and population (NBS; WPP) as the economic indicator. The

saturated vehicle stock per 1000 people was first derived from previous studies (Huo and Wang, 2012) and then adjusted by the maximal vehicle stock per 1000 people calculated using statistical data. The combined vehicle statistical database was used to estimate parameters (α and β) of the Gompertz function at the country level. For countries whose R square (R^2) of the country-level regression was less than 0.5, regional or global α and β regression parameters were used instead (Zheng et al., 2012).

As the verification of the vehicle stock modeled by the Gompertz function, we compared them with the statistical vehicle stock for countries in years when statistics were available. The relative deviation ratios in countries that own top 85% of global vehicles stock were between -28% and 25.6%, ranges of the relative deviation in rest countries were a bit larger due to the limited availability of statistics. Figure 2(a) and Figure S3 show the comparison in 2015, a year with more statistical data. The deviation of the modeled vehicle stock from the statistics in most countries was less than $\pm 25\%$, especially in the United States, countries in the European Union, China, and India. The relatively good consistency between the modeled and statistical vehicle stock indicates the relatively high reliability of this model. Therefore, a long-term integrated vehicle stock database (1970-2020) was constructed by constraining the modeled vehicle stock by the combined vehicle statistical database.




Figure 2: Verification of the modeled vehicle stock in United States, the European Union, China, and India (a) and the age distribution for PLDVs (b) in 2015.

2.3 Modeling the age distribution of vehicle stock

Then, the age distribution of the stock was modeled using the dynamic balanced function with the integrated vehicle stock database set up in the first step and a combined vehicle sale statistical database. Similar to the combination of vehicle stock statistics, OICA data were used as the default vehicle sale database with WRS data as a supplement after comparison, and local statistics (ACEA; CEIC; EC; JAMA; MEIC; NBS; TEDB) were also involved to correct the default database. Limited by the temporal coverage of the statistical data, vehicle sales were not available for most countries before 2005. Therefore,

the newly registered vehicles for missing years was back-calculated with the dynamic balanced function, in which the vehicle stock from the previous step and survival rates derived from available studies and reports (Huo and Wang 2012; Yan et al., 2011; Yan et al., 2014; Zheng et al., 2014) were inputs. Here we marked 231 countries into two types: focus countries and broader regions (Table S1). 20 countries owning the top 75% of global vehicles were marked as focus countries, for which the dynamic balanced function was built at country level. The remaining 211 countries were marked as broader regions and further combined into 8 regions according to the roadmap region definition (ICCT 2012). In each broader region, data in a representative country, which has most abundant statistics with region, was used to build the dynamic balanced function and the age distribution in this country was assumed to be able to represent that in other countries belonging to the same region. The age distribution in this study was not simulated for MCs due to the limitation of data availability, and we assumed that they shared the same age distribution of PLDVs.

To verify the age distribution modeled by the dynamic balanced function, relative deviation between the simulated vehicle stock based on newly registered vehicles and survival rates and the vehicle stock in the first step was used as the validation indicator. Except for several years in Argentina and Thailand, the relative deviation ratios of light-duty vehicles during 1970-2020 ranges from -30.9% to 30.8%, heavy-duty vehicles had larger relative deviation ratios which were between -36.5% and 34.9%. Taking 2015 as an example, the relative deviation ratios in most countries were less than $\pm 30\%$ (Figure 2(b) and Figure S4). The relatively good consistency between the vehicle stock and simulation indicated that the dynamic balance function set up in this study could well model the entry of newly registered vehicles and the retirement of existing vehicles and the estimated age distribution was reliable.

2.4 Estimates of fuel consumption

In the third step, we estimated the initial vehicular fuel consumption based on outputs from the first two steps and parameters including the annual average kilometers traveled (VKT), fuel structure, and fuel economy. Then the initial vehicular fuel consumption was constrained with energy statistics from World Energy Statistics (IEA¹) at country level, which was finally used in CO₂ estimation. VKT, fuel structure, and fuel economy are rarely available in global statistics annually, this study used fleet-average data, which were estimated based on vehicle-kilometers, the vehicle stock, vehicle-kilometer energy intensity, and fuel consumption by category in energy efficiency statistics (IEA²). These indexes for 39 countries

(accounting for 43%-73% of the global vehicle stock) during the 2000-2018 period can be found in energy efficiency statistics. For countries that were not covered in energy efficiency statistics, the regional or global mean VKT, fuel structure, and fuel economy were used. For missing years, we assumed that the values of these three parameters were similar to those of the adjacent year. There are few local statistics or studies that evaluate the VKT, fuel structure, and fuel economy; therefore, these parameters were supplemented and revised only for the United States, Europe, China, and Japan using local statistics or studies (AECA; IEA³; JAMA; MEIC; TEDB; TRACCS).

As the validation of fuel consumption, the initial vehicular fuel consumption was compared to energy statistics by fuel type (Figure S5). The range of relative deviation ratios of gasoline, diesel, NG, and LPG was -23% to 3%, -19% to 9%, -22% to 34%, and -39% to 14%, respectively. As CO₂ is not directly emitted as exhaust by electrical vehicles whether they were running, starting or parking, electricity was not considered in the estimation of vehicular fuel consumption in this study. The consistency of the simulation with statistics ensured the feasibility of constraining the modeled fuel consumption by statistics.

2.5 Estimates of CO₂ emissions and uncertainty assessment

Finally, vehicular CO₂ emissions were estimated using the constrained vehicular fuel consumption from previous step and a combined CO₂ emission factor database in which emission factors from the Intergovernmental Panel on Climate Change (IPCC) were used as the default emission factors, and local studies (EEA; Shan et al., 2018) were used as supplements and amendments. As the CO₂ emission factor is influenced mainly by the fuel type and country, the estimation of CO₂ emissions would not be interfered with by the simplified assumption for MCs in modelling the age distribution.

Following the method in Crippa et al. (2018) and Crippa et al. (2019), the corresponding uncertainty (σ) of CO₂ emissions from on-road vehicles in year y for a given country c is calculated as following:

$$\sigma_{Emis_{c,y}} = \sqrt{\sum_f \left(\sigma_{AD_{c,y,f}}^2 + \sigma_{EF_{c,f}}^2 \right) \times \left(Emis_{c,y,f} / Emis_{c,y} \right)^2} \quad (6)$$

where σ_{AD} and σ_{EF} are the uncertainties (%) of the activity data (the constrained fuel consumption of on-road vehicles) and CO₂ emission factors. Based on assumption of lognormal distribution of the calculated uncertainties (Bond et al., 2004), we evaluated the upper and lower range of CO₂ estimate by multiplying and dividing the base emissions in this study by $(1 + \sigma)$, respectively (Crippa et al., 2018).

As CO₂ uncertainty can vary significantly among countries (Marland et al., 1999; Olivier et al., 2014) and the primary source of uncertainty of the CO₂ estimate from on-road vehicles is the activity data rather than emission factors (GPG 2000), the main step in CO₂ uncertainty assessment is to evaluate the uncertainty of national activity data. In this study, 231 countries were divided into several groups (Table S1) in the uncertainty assessment in accordance with IPCC tiered approach and EDGAR (Janssens-Maenhout et al., 2019). Here we assume that countries belonging to the OECD in 1990 (OECD90) have the lowest uncertainties in their fuel consumption data because they were economically stable and would have a good statistical infrastructure. On the same line, fuel consumption data in countries with Economies in Transition of 1990 (EIT90) is more uncertain than that of OECD90 but less than that from the other remaining non-Annex I countries. Exceptions to the country grouping are made for Australia, Canada, China, India, Japan, Russia, Ukraine, United States, and countries belonging to the 15 member countries of European Union (EU15) whose uncertainty values of fuel consumption data were obtained from Olivier et al. (2016) and Hong et al. (2017). Uncertainty values for CO₂ emission factors were retrieved from EEA.

Table S4 shows the corresponding uncertainty of CO₂ emissions at both global and regional level during 1970-2020 on basis of Equation 6. The uncertainty in the global on-road CO₂ emissions is estimated to range from -7.2% to 8.1%, which is close to the expert judgement suggested value (approximately $\pm 5\%$) in GPG (2000). Because sufficient local data was used in the CO₂ estimation, United States and European Union have the lowest uncertainty in the range of -3.8% to 4.0% and -2.9% to 3.0%, respectively. India also has relatively low uncertainty that varies between -4.7% and 5.0% because of the low uncertainty derived from Janssens-Maenhout et al. (2019) in which India is classified as countries with well-developed statistical systems. Due to the less-developed statistical systems, Latin Am. + Canada and Middle East + Africa have the largest uncertainty, which range from -12.3% to 14.6% and -15.4% to 18.3%, respectively. Hong et al. (2017) found that the apparent uncertainties in oil consumption statistics in China during 1996-2003 were relatively large with an average apparent uncertainty ratio of 15.8%, which led to the relatively larger uncertainty in China's on-road CO₂ emissions with the range of -12.6% to 14.4%. It could also be found that uncertainties at regional level decreased over time with the development of statistical systems in more countries. But uncertainty in global on-road CO₂ emissions slightly increased during 1970-2020 due to the growing contribution of regions with larger uncertainty to the global total CO₂ emissions.

3 Results

3.1 Evolution of the global vehicle stock, 1970-2020

The global vehicle stock continuously increased from 0.3 billion in 1970 to 2.3 billion in 2020, and there is both consistency and variety between countries in terms of the distributions of vehicles and fuel types (Figures 3 and S7). In 1970, PLDVs were the major vehicle type in United States (83%) and the European Union (88%) but had relatively low proportions in China (23%) and India (5%). The high proportion of PLDVs in the United States and the European Union, as well as the dominant position of these two regions in terms of the global vehicle stock (Figure S6), led to more than 70% of global vehicles being PLDVs in 1970. The proportion of PDLVs in China significantly increased and reached 68% in 2020 and have replaced MCs to become the dominant vehicle type. Although the stock of PLDVs in India also increased substantially during the 1970-2020 period, MCs with the proportion of 78% the vehicle stock in 2020 were still the most frequently used vehicles in India, benefiting by the local warm climate. The majority of vehicles in the European Union in 2020 were still PLDVs, for which the proportion was 79%, but the dominant vehicle type in United States has changed from PLDVs to CLDVs and CLDVs accounted for 50% of the local vehicle stock. As the dominant position of developed countries in global vehicle stock replaced by developing countries during the 1970-2020 period (Figure S6), the share of MCs in the global vehicle stock increased accordingly to 32%, and the proportion of PLDVs decreased to 50% in 2020.

Unlike the changes in the vehicle-type distribution during the 1970-2020 period, the fuel structure of the vehicle stock was consistent in most regions. Currently, the majority of the vehicle stock worldwide still consists of gasoline and diesel vehicles, which together accounted for 98% of the global vehicle stock in 2020. Gasoline was the major fuel type for vehicles in most countries from 1970 to 2020, but the dieselization of PLDVs in regions such as the European Union (Figure S10) led to a larger proportion of diesel vehicles in the local vehicle stock. For example, the share of diesel vehicles in the European Union increased from 29% in 1970 to 43% in 2020. Although the share of electrical vehicles in the vehicle stock was still much lower than that of gasoline and diesel vehicles, the stock of global electrical PLDVs has reached 10.2 million, and in this regard, the growth has been the fastest in the last eight years.




Figure 3: Trends in vehicle ownership from 1970 to 2020.

3.2 CO₂ emissions from global on-road vehicles

Global CO₂ emissions from on-road vehicles continued to increase overall from 1.7 Gt in 1970 to 5.4 Gt in 2020 (Figure 4). Profiting from the integrated global vehicle database developed in this study, we further analyzed the vehicle- and fuel type-specific characteristics of CO₂ emissions from global on-road vehicles. On-road CO₂ emissions were concentrated in specific vehicle and fuel types throughout the period. From 1970 to 2020, almost all of global CO₂ emissions from on-road vehicles came from gasoline and diesel vehicles due to their dominant proportion in the vehicle stock (Figure S10). In 1970, 78% and 21.5% of global on-road CO₂ emissions were exhausted from gasoline and diesel vehicles, respectively, and in 2020, these emissions together accounted for 96% of global on-road CO₂ emissions; only the ranking of the contributions changed. With continuous dieselization during the 1970-2020 period (Figure S10), the contribution of diesel vehicles to global on-road CO₂ emissions increased to 47% in 2020. Although CO₂ emissions from vehicles using other fuels (here, NG and LPG) continued to grow during the 1970-2020 period, their proportions were still quite slight compared to those of gasoline and diesel vehicles.

PLDVs, accounting for the largest share in the global vehicle stock, were also the main source of global on-road CO₂ emissions and contributed more than 47% of global CO₂ emissions from on-road

vehicles during the 1970-2020 period. Although MCs accounted for the second largest share in the global vehicle stock, CO₂ emissions from MCs were not comparable to those from PLDVs. In 2020, proportion of PLDVs and MCs in the global vehicle stock was 50% and 32%, respectively, and their CO₂ emissions were 2.6 Gt and 0.3 Gt, respectively, which accounted for 48% and 5% of global on-road CO₂ emissions, respectively. In contrast, trucks with a fairly low share in the global vehicle stock contributed the second largest share of global on-road CO₂ emissions. During the 1970-2020 period, trucks accounted for less than 5% of the global vehicle stock but exhausted 17% of global on-road CO₂ emissions in 1970, and their contribution increased to 22% in 2020. As most PLDVs are gasoline vehicles and the majority of trucks are powered by diesel, gasoline PLDVs and diesel trucks are among the top 2 vehicle- and fuel type-specific contributors to global on-road CO₂ emissions. In 2020, the CO₂ emissions from gasoline PLDVs and diesel trucks were 1.8 Gt and 1.1 Gt, respectively, accounting for 33% and 20% of global on-road CO₂ emissions, respectively.




Figure 4: Global CO₂ emissions from 1970 to 2020 by vehicle and fuel type. The panels are organized by fuel type (rows) and vehicle type (columns).

Figure 5 shows the geographical distribution of the two largest contributors to global on-road CO₂ emissions in 2020, namely, gasoline PLDVs and diesel trucks. Global on-road CO₂ emissions were highly concentrated in several countries. In 2020, the top 10 countries contributed 69% and 71% of global CO₂ emissions exhausted from gasoline PLDVs and diesel trucks, respectively. The United States was still the largest contributor to global CO₂ emissions from both gasoline PLDVs and diesel trucks, whose contributions were up to 25% and 28%, respectively. With the continuous improvement in China's economic development, China became the leading market for global vehicles in 2020 (Figure S6) and accounted for 18% and 19% of CO₂ emissions from global gasoline PLDVs and diesel trucks, respectively. Although growth in on-road CO₂ emissions in developed countries slowed down after 2000

(Figure S8), the contributions of gasoline PLDVs and diesel trucks in developed countries were still greater than those in developing countries, especially for gasoline PLDVs. For example, the ownership of gasoline PLDVs in Canada and India was relatively close in 2020, at 22.5 and 21.2 million, respectively, but the CO₂ emissions from gasoline PLDVs in Canada were 83.5 Mt, which is three times greater than that in India.




Figure 5: Maps of on-road CO₂ emissions from the top 2 contributors worldwide: (a) gasoline PLDVs and (b) diesel trucks.

We further analyzed the influence of shifts in the fuel-type distribution of vehicle ownership (Figure S10) on the fuel structure of on-road CO₂ emissions (Figure 6 and Figure S11). In 1970, CO₂ emissions from PLDVs were mainly exhausted from gasoline vehicles, as the majority of PLDVs in most regions were powered by gasoline, and diesel vehicles exhausted only 7% of CO₂ emissions from PLDVs worldwide. In 2020, gasoline vehicles were still the dominant contributor to CO₂ emissions from PLDVs in the United States and China, but the contribution of diesel vehicles increased significantly in the European Union and India, which accounted for 61% and 50% of local CO₂ emissions from PLDVs, respectively. Influenced by the dieselization of PLDVs in regions such as the European Union and India, the contribution of diesel vehicles to CO₂ emissions from PLDVs in 2020 also increased to 28%. For

CLDVs, the contribution of diesel vehicles was more than 50% in the European Union, China, and India, but in the remaining regions, CO₂ emissions were still mainly from gasoline vehicles. Buses and trucks were also dieselized during the 1970-2020 period, and diesel vehicles have become the dominant contributor to CO₂ emissions from buses and trucks both regionally and globally. Therefore, controlling emissions from diesel vehicles, especially buses and trucks, holds great significance for reducing global on-road CO₂ emissions.




Figure 6: Transition of diesel vehicles' contribution to CO₂ emissions.

3.3 Age distribution of CO₂ emissions

On the basis of the fleet turnover emission model built in this study, the age distribution of global on-road CO₂ emissions was estimated and analyzed (Figure 7). The contribution of old vehicles (those that survived more than 15 years) to CO₂ emissions was relatively low, regardless of whether they were light-duty or heavy-duty vehicles. In 1970, old vehicles contributed 4% and 6% of CO₂ emissions from light-duty and heavy-duty vehicles, respectively. Although the contribution of old vehicles to CO₂ emissions increased, they still contributed only approximately 10% of CO₂ emissions from both light-duty and heavy-duty vehicles in 2020. As emissions of air pollutants such as particulate matter (PM) may increase with age because of degradation in engine performance and air pollution control equipment (Yan et al., 2011), the contributions of old vehicles to emissions of air pollutants could be much greater than those

of CO₂. Therefore, controlling old vehicles may not be significant in mitigating CO₂ emissions but could lead to effective air pollutant emission coreductions.

Global CO₂ emissions from vehicles of all ages were mainly contributed by developed countries, such as the United States and countries in the European Union before 2000, as these countries owned the majority of global vehicles during that period. After 2000, the contributions of vehicles in developing countries such as China and India to global on-road CO₂ emissions increased significantly, especially for CO₂ emissions from vehicles younger than ten years. Taking CO₂ emissions from light-duty vehicles aged 0-1 as an example, the proportion of these vehicles in China increased from 1% in 1970 to 16% in 2020, while the proportion of these vehicles in the United States decreased from 44% in 1970 to 23% in 2020. CO₂ emissions from old vehicles in 2020 were still mainly exhausted by vehicles in developed countries such as the United States and countries in the European Union, which is related to the longer lifetimes and earlier development of vehicles in these countries. For example, old vehicles in the United States contributed nearly half of the CO₂ emissions exhausted from old light-duty vehicles worldwide in 2020.




Figure 7: Shares of CO₂ emissions by vehicle age. In each panel, the bars from left to right show the proportions of the world, the United States (US), the European Union (EU), China, and India accounted for by vehicles in the vehicle age categories. The panels are organized by year (rows) and vehicle type (columns).

4 Data availability

The fuel-, vehicle type-, and age-specific CO₂ emission data presented herein cover the period from 1970 to 2020 at the country level. The data are available as open data at <https://doi.org/10.6084/m9.figshare.24548008> (Yan et al., 2023).

5 Conclusions

Our study constructed a fuel-, vehicle type-, and age-specific CO₂ emission inventory from 1970 to 2020 of global on-road vehicles covering 231 countries, five types of fuel, and five types of vehicles. In this model, the best available statistics on the vehicle stock and sales were used to model the vehicle stock via the Gompertz function as well as the age distribution based on the dynamic balanced relationship between the vehicle stock and vehicle sales. Statistical fuel consumption was used to constrain the estimated vehicular fuel consumption at the country level, and emission factors from both the IPCC and local studies were used to estimate CO₂ emissions. On the basis of our CO₂ emission inventory with detailed information, the evolution of the global vehicle stock over 50 years was analyzed, the dominant emission contributors by vehicle and fuel type were identified, and the age distribution of on-road CO₂ emissions was also characterized. We found that trucks accounted for less than 5% of global vehicle ownership but represented more than 20% of on-road CO₂ emissions in 2020. The contribution of diesel vehicles to global on-road CO₂ emissions doubled during the 1970-2020 period, driven by the shift in the fuel-type distribution of vehicle ownership. The proportion of CO₂ emissions from vehicles in developing countries such as China and India in terms of global emissions from newly registered vehicles significantly increased after 2000, but global CO₂ emissions from vehicles that survived more than 15 years in 2020 still originated mainly from developed countries such as the United States and countries in the European Union.

The fleet turnover model built in this study could also be used for estimating global on-road emissions of air pollutants, which are more significantly influenced by the vehicle-type distribution, fuel structure, and age distribution of the fleet. However, these fuel-, vehicle type-, and age-specific characteristics have not yet been discussed in existing studies. In the future, our model could help improve the global emission inventory of air pollutants from on-road vehicles and further support analyses of coreductions in CO₂ and air pollutant emissions from global on-road vehicles as well as the

potential air quality and climate cobenefits. In addition to the uncertainty quantification for our CO₂ emission data, we further verified the reliability of CO₂ emissions in this study by comparing them to those of other widely used global, regional, and national emission inventories in which long-term CO₂ emissions are available (Figure S12). The CO₂ emissions in this study not only exhibited good consistency with other global emission inventories at the global scale but also were more similar to local emissions than those in other global or regional emission inventories at the country and regional levels.

Supplement. The data related to figures in this article is available in the supplementary file Figures.zip.

Author contributions. LY collected the data, developed the fleet turnover model, and constructed the database of fuel-, vehicle type-, and age-specific CO₂ emissions from global on-road vehicles during the 1970-2020 period. LY and QZ discussed the expansion of the database. LY wrote the paper with the help of all the coauthors.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Earth System Science Data.

Acknowledgments. This work was supported by the National Natural Science Foundation of China (41921005, 42130708, and 72140003).

References

- Amstel, A. v., Olivier, J., and Janssen, L.: Analysis of differences between national inventories and an Emissions Database for Global Atmospheric Research (EDGAR), *Environ. Sci. Policy*, 2, 275-293, [https://doi.org/10.1016/S1462-9011\(99\)00019-2](https://doi.org/10.1016/S1462-9011(99)00019-2), 1999.
- Beven, K.: Facets of uncertainty: epistemic uncertainty, non-stationarity, likelihood, hypothesis testing, and communication, *Hydrolog. Sci. J.*, 61, 1652–1665, <https://doi.org/10.1080/02626667.2015.1031761>, 2016.
- Bhalla, K., Brauer, M., Burnett, R., Cohen, A. R., Freedman, G., Leach-Kemon, K., Murray, C. J. L., Shahraz, S., Shotten, M. S.: Transport for Health: The Global Burden of Disease from Motorized Road Transport. <http://documents.worldbank.org/curated/en/984261468327002120/Transport-for-health-the->

419 global-burden-of-disease-from-motorized-road-transport (accessed 04 January 2024) (Global Road
420 Safety Facility, Institute for Health Metrics and Evaluation, The World Bank, 2014).

421 Boden, T. A., Marland, G., Andres, R. J.: Global, Regional, and National Fossil-Fuel CO₂ Emissions,
422 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of
423 Energy, Oak Ridge, Tenn., USA, https://doi.org/10.3334/CDIAC/00001_V2016, 2016.

424 Boden, T. A., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO₂ Emissions,
425 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of
426 Energy, Oak Ridge, Tenn., USA, https://doi.org/10.3334/CDIAC/00001_V2017, 2017.

427 Bond, T. C.: A technology-based global inventory of black and organic carbon emissions from
428 combustion. *Journal of Geophysical Research*, 109, (D14), <https://doi.org/10.1029/2003JD003697>, 2004.

429 CEIC: <https://insights.ceicdata.com> (accessed 04 January 2024).

430 Crippa, M., Janssens-Maenhout, G., Guizzardi, D., Van Dingenen, R., Dentener, F.: Contribution and
431 uncertainty of sectorial and regional emissions to regional and global PM_{2.5} health impacts. *Atmospheric*
432 *Chemistry and Physics*, 19(7), 5165-5186, <https://doi.org/10.5194/acp-19-5165-2019>, 2019.

433 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S.,
434 Doering, U., Olivier, J. G. J., Pagliari, V., Janssens-Maenhout, G.: Gridded emissions of air pollutants
435 for the period 1970–2012 within EDGAR v4.3.2. *Earth System Science Data*, 10, (4), 1987-2013,
436 <https://doi.org/10.5194/essd-10-1987-2018>, 2018.

437 Crippa, M., Janssens-Maenhout, G., Guizzardi, D., Galmarini, S.: EU effect: Exporting emission
438 standards for vehicles through the global market economy. *Journal of Environ Manage*, 183, 959-971,
439 <http://dx.doi.org/10.1016/j.jenvman.2016.09.068>, 2016.

440 Dargay J, Gately D, Sommer M. Vehicle Ownership and Income Growth, Worldwide: 1960-2030. *The*
441 *Energy Journal*, 28(4), <https://doi.org/10.5547/ISSN0195-6574-EJ-VOL28-NO4-7>, 2007.

442 Dargay J, Gately D. Income's effect on car and vehicle ownership, worldwide: 1960 –2015.
443 *Transportation Research Part A. Policy and Practice*, 33(2), [https://doi.org/10.1016/S0965-](https://doi.org/10.1016/S0965-8564(98)00026-3)
444 [8564\(98\)00026-3](https://doi.org/10.1016/S0965-8564(98)00026-3), 1999.

445 Davis, S. J., S. Lewis, N. S., Shaner, M. R., Aggarwal, S., Arent, D. J., Azevedo, I., Benson, S. M.,
446 Bradley, T. H., Brouwer, J., Chiang, Y. M., Clack, C. T. M., Cohen, A., Doig, S. J., Edmonds, J., Fennell,
447 P. S., Field, C. B., Hannegan, B., Hodge, B. M., Hoffert, M. I., Ingersoll, E., Jaramillo, P., Lackner, K.
448 S., Mach, K. J., Mastrandrea, M. D., Ogden, J. M., Peterson, P. F., Sanchez, D. L., Sperling, D., Stagner,

449 J., Trancik, J. E., Yang, C. J., Caldeira, K. Net-zero emissions energy systems. *Science*, 360(6496),
 450 <https://doi.org/10.1126/science/aas9793>, 2018.

451 European Automobile Manufacturers Association (ACEA): Vehicles in use Europe 2017,
 452 <https://www.acea.auto/> (accessed 04 January 2024)

453 European Commission (EC): <https://ec.europa.eu/> (accessed 04 January 2024)

454 European Environment Agency (EEA): EMEP/EEA air pollutant emission inventory guidebook 2019,
 455 http://efdb.apps.eea.europa.eu/?source=%7B%22query%22%3A%7B%22match_all%22%3A%7B%7D%2C%22display_type%22%3A%22tabular%22%7D (accessed 04 January 2024)

456 GPG (2000): Good Practice Guidance and Uncertainty Management in National Greenhouse Gas
 457 Inventories. Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L.,
 458 Hoppaus R., Martinsen T., Meijer J., Miwa K., Tanabe K. (Eds). Intergovernmental Panel on Climate
 459 Change (IPCC), IPCC/OECD/IEA/IGES, Hayama, Japan. [https://www.ipcc-](https://www.ipcc-nggip.iges.or.jp/public/gp/english/)
 460 [nggip.iges.or.jp/public/gp/english/](https://www.ipcc-nggip.iges.or.jp/public/gp/english/) (accessed 04 February 2024)

461 Hoesly, R. M., Smith, S. J., Feng, L.; Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J.,
 462 Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-i., Li, M., Liu,
 463 L., Lu, Z., Moura, M. C. P., Rourke, P. R.; Zhang, Q.: Historical (1750–2014) anthropogenic emissions
 464 of reactive gases and aerosols from the Community Emissions Data System (CEDS). *Geoscientific*
 465 *Model Development*, 11, (1), 369-408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

466 Hong, C., Zhang, Q., He, K., Guan, D., Li, M., Liu, F., Zheng, B.: Variations of China's emission
 467 estimates: response to uncertainties in energy statistics. *Atmospheric Chemistry and Physics*, 17(2):
 468 1227-1239, <https://doi.org/10.5194/acp-17-1227-2017>, 2017.

469 Huo, H., Wang, M.: Modeling future vehicle sales and stock in China. *Energy Policy*, 43: 17-29,
 470 <https://doi.org/10.1016/j.enpol.2011.09.063>, 2012.

471 Intergovernmental Panel on Climate Change (IPCC): 2006 IPCC Guidelines for National Greenhouse
 472 Gas Inventories, <https://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (accessed 04 January 2024)

473 International Energy Agency (IEA1): Energy Statistics and Balances of Non-OECD Countries, 1960–
 474 2020, Paris, 2022b

475 International Energy Agency (IEA1): Energy Statistics and Balances of OECD Countries, 1960–2020,
 476 Paris, 2022a

477 International Energy Agency (IEA2): Energy Efficiency Statistics 2019 Edition, Paris, 2019

479 International Energy Agency (IEA3): Global EV Outlook 2023, [https://www.iea.org/reports/global-ev-](https://www.iea.org/reports/global-ev-outlook-2023)
480 outlook-2023, (accessed 04 January 2024)

481 International Road Federation (IRF): IRF World Road Statistics (WRS) 2021 Edition,
482 <https://worldroadstatistics.org/> (accessed 04 January 2024)

483 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi,
484 P., Pagliari, V., Olivier, J. G. J., Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu,
485 A. M. R., Solazzo, E., Oreggioni, G. D.: EDGAR v4.3.2 Global Atlas of the three major greenhouse gas
486 emissions for the period 1970–2012. *Earth System Science Data*, 11, (3), 959–1002,
487 <https://doi.org/10.5194/essd-11-959-2019>, 2019.

488 Japan Automobile Manufacturers Association (JAMA): <https://www.jama.org/>, (accessed 04 January
489 2024).

490 Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., Schöpp, W.:
491 Global anthropogenic emissions of particulate matter including black carbon. *Atmospheric Chemistry*
492 *and Physics*, 17(14), 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.

493 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., Pozzer, A.: The contribution of outdoor air pollution
494 sources to premature mortality on a global scale. *Nature*, 525, 367–371,
495 <https://doi.org/10.1038/nature16371>, 2015.

496 Marland, G., Brenkert, A., and Olivier, J.: CO₂ from fossil fuel burning: A comparison of ORNL and
497 EDGAR estimates of national emissions, *Environ. Sci. Policy*, 2, 265–274, 1999.

498 Matthews, H. D., Caldeira, K.: Stabilizing climate requires near-zero emissions. *Geophys. Res. Lett.*, 35,
499 L04705, <https://doi.org/10.1029/2007GL032388>, 2008.

500 Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C., Frieler, K., Knutti, R., Frame, D. J.,
501 Allen, M. R.: Greenhouse-gas emission targets for limiting global warming to 2 °C. *Nature*, 458, 1158–
502 1162, <https://doi.org/10.1038/nature08017>, 2009.

503 Ministry of Statistics and Programme Implementation (MOSPI): <https://mospi.gov.in/>, (accessed 04
504 January 2024).

505 Multi-resolution Emission Inventory model for Climate and air pollution research (MEIC):
506 <http://meicmodel.org.cn/>, (accessed 04 January 2024).

507 National Bureau of Statistics of China (NBS): <https://data.stats.gov.cn/>, (accessed 04 January 2024).

508 Hähne, N., Elzen, M. D., Rogelj, J., Metz, B., Fransen, T., Kuramochi, T., Olhoff, A., Alcamo, J.,
 509 Winkler, H., Fu, S., Schaeffer, M., Schaeffer, R., Peters, G. P., Maxwell, S., Dubash, N. K.: Emissions:
 510 world has four times the work or one-third of the time, *Nature*, 579, 25-28,
 511 <https://doi.org/10.1038/d41586-020-00571-x>, 2020.

512 Oda, T., Maksyutov, S., Andres, R. J.: The Open-source Data Inventory for Anthropogenic CO₂, version
 513 2016 (ODIAC2016): a global monthly fossil fuel CO₂ gridded emissions data product for tracer transport
 514 simulations and surface flux inversions. *Earth System Science Data*, 10(1), 87-107,
 515 <https://doi.org/10.5194/essd-10-87-2018>, 2018.

516 Olivier, J. G. J., Janssens-Maenhout, G., Muntean, M., and Peters, J. A. H. W.: Trends in global CO₂
 517 emissions: 2014 report, European Commission – PBL Netherlands Environmental Assessment Agency,
 518 The Hague, JRC93171/PBL1490 report, ISBN 978-94-91506-87-1, 2014.

519 Olivier, J. G. J., Janssens-Maenhout, G., Muntean, M., and Peters, J. A. H. W.: Trends in global CO₂
 520 emissions: 2016 Report, no. 2315, PBL Netherlands Environmental Assessment Agency, The Hague,
 521 2016.

522 Organisation Internationale des Constructeurs d'Auto (OICA): World vehicle in use: 2005-2015.
 523 <https://www.oica.net/category/vehicles-in-use/>, (accessed 04 January 2024).

524 Rogelj, J., D. Shindell, K. Jiang, S. Fifita, P. Forster, V. Ginzburg, C. Handa, H. Kheshgi, S. Kobayashi,
 525 E. Kriegler, L. Mundaca, R. S. Ą́rian, and M.V. Vilari ño,: Mitigation Pathways Compatible with 1.5 °C
 526 in the Context of Sustainable Development. In: Global Warming of 1.5 °C. An IPCC Special Report on
 527 the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas
 528 emission pathways, in the context of strengthening the global response to the threat of climate change,
 529 sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner,
 530 D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péán, R. Pidcock, S. Connors, J.B.R.
 531 Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)].
 532 Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 93-174,
 533 <https://doi.org/10.1017/9781009157940.004>, 2018.

534 Shan, Y., Guan, D., Zheng, H., Ou, J., Li, Y., Meng, J., Mi, Z., Liu, Z., Zhang, Q.: China CO₂ emission
 535 accounts 1997–2015. *Scientific Data*, 5(1), 170201, <https://doi.org/10.1038/sdata.2017.201>, 2018

536 Shindell, D., Faluvegi, G., Walsh, M., Anenberg, S. C., Milly, G.: Climate, health, agricultural and
 537 economic impacts of tighter vehicle-emission standards. *Nat. Clim. Change*, 1, 59–66, [https://doi.org/](https://doi.org/10.1038/nclimate1066)
 538 10.1038/nclimate1066, 2011.

539 Silva, R., Adelman, Z., Fry, M. M., West, J. J.: The impact of individual anthropogenic emissions sectors
 540 on the global burden of human mortality due to ambient air pollution. *Environ. Health Perspect*, 124,
 541 1776–1784, <https://doi.org/10.1289/EHP177>, 2016.

542 The International Council on Clean Transportation (ICCT): Global Transportation Roadmap Model
 543 Documentation and User Guide, 2012

544 Transport data collection supporting the quantitative analysis of measures relating to transport and
 545 climate change (TRACCS): <https://traccs.emisia.com/>, (accessed 04 January 2024).

546 Transportation Energy Data Book (TEDB): <https://tedb.ornl.gov/data>, (accessed 04 January 2024).

547 UNdata: <http://data.un.org/>, (accessed 04 January 2024).

548 Unger, N., Bond, T. C., Wang, J. S., Koch, D. M., Menon, S., Shindell, D. T., Bauer, S.: Attribution of
 549 climate forcing to economic sectors. *Proc. Natl Acad. Sci.*, 107, 3382–3387,
 550 <https://doi.org/10.1073/pnas.0906548107>, 2010.

551 Wang, R., Tao, S., Ciais, P., Shen, H. Z., Huang, Y., Chen, H., Shen, G. F., Wang, B., Li, W., Zhang, Y.
 552 Y., Lu, Y., Zhu, D., Chen, Y. C., Liu, X. P., Wang, W. T., Wang, X. L., Liu, W. X., Li, B. G., Piao, S.
 553 L.: High-resolution mapping of combustion processes and implications for CO₂ emissions. *Atmospheric*
 554 *Chemistry and Physics*, 13(10), 5189–5203, <https://doi.org/10.5194/acp-13-5189-2013>, 2013.

555 World Bank (WB): <https://data.worldbank.org/>, (accessed 04 January 2024).

556 World Population Prospects 2022 (WPP): <https://population.un.org/wpp/>, (accessed 04 January 2024).

557 Yan, F., Winijkul, E., Bond, T. C., Streets, D. G.: Global emission projections of particulate matter (PM):
 558 II. Uncertainty analyses of on-road vehicle exhaust emissions. *Atmospheric Environment*, 87, 189–199,
 559 <http://dx.doi.org/10.1016/j.atmosenv.2014.01.045>, 2014.

560 Yan, F., Winijkul, E., Jung, S., Bond, T. C., Streets, D. G.: Global emission projections of particulate
 561 matter (PM): I. Exhaust emissions from on-road vehicles. *Atmospheric Environment*, 45(28), 4830–4844,
 562 <https://doi.org/10.1016/j.atmosenv.2011.06.018>, 2011.

563 Yan, L., Zhang, Q., He, K. B.: CO₂ emissions from global on-road vehicles by vehicle- and fuel- type
 564 for each country during 1970–2020. <https://doi.org/10.6084/m9.figshare.24548008>.

565 Zheng, B., Huo, H., Zhang, Q., Yao, Z. L., Wang, X. T., Yang, X. F., Liu, H., He, K. B.: High-resolution
566 mapping of vehicle emissions in China in 2008. *Atmospheric Chemistry and Physics*, 14(18), 9787-9805,
567 <https://doi.org/10.5194/acp-14-9787-2014>, 2014